

Purpose: To observe and list biotic factors that affect a given ecosystem.

- Summary: In this exercise, students will identify and observe biotic factors in a riparian ecosystem, which is the green strip of vegetation alongside a waterbody; they will measure the types of vegetation at the water's edge, the function of plants as ground cover and canopy cover, and observe the wildlife in the area.
- Background: Riparian areas are the transition zone between aquatic and terrestrial systems. The plants in this zone depend on high water tables and flooding patterns associated with the aquatic systems. Riparian zones cover a very small area in a watershed, but are extremely important.

Riparian Zone Functions:

- Bank stability to resist erosion
- Habitat for a diverse community of plants and animals
- Canopy cover which provides shading
- Organic materials drop from canopy cover into waterbodies
- Soils soak up water from runoff and prevent flooding
- Healthy riparian areas protect the land from flooding, and provide storage for a sustained summer flow

For more background information, see:

• The Riparian Zone section of the Utah Stream Team Manual which defines a riparian zone and discusses how it would change due to natural and human influences, why the riparian zone is important in an aquatic ecosystem, and how to measure and interpret the results.

• For information on Utah's wildlife, see the Utah Division of Wildlife Resources Data Center website – http://dwrcdc.nr.utah.gov/ucdc Duration: Classroom 20 minutes Outdoors 50 min.


Setting: Classroom Outdoors

Link to the Utah Core Curriculum: Earth Systems– 9thgrade Standard II-2b Standard II-3a

L	0	's	:

1 a, f, g, l, j 2 b, c, e 3 a, c, d 4 a-e 6 a-d, h

USU Water Quality Extension - Stream Side Science





• Greenline - they will measure the type of vegetation that grows closest to the water's edge. This is an indication of the bank stability. See Further Discussion question number 1 for more information.

• Ground Cover - they will record the width of the zone in which riparian plants grow.

• Canopy Cover - they will measure the amount of shade the riparian plants provide.

• Wildlife Signs - they will identify animals and signs of animal activity.

Because there are four sets of measurements, we strongly recommend reviewing the actual measuring procedures with the class before going into the field.

Field Activity:

1. Divide your students into groups of no more than six students.

2. Assign each group with a measurement (greenline, ground or canopy cover or wildlife signs) and provide them with the appropriate materials.

Greenline Group

Flagging Measuring tape Greenline sampling instructions Greenline worksheet

Canopy Cover Group

Ocular tube Measuring tape Canopy cover sampling instructions Canopy cover worksheet

Ground Cover Group

Measuring tape Ground cover sampling instructions Ground cover worksheet

Safety First! Always consider

safety factors when working near water.

Wildlife Group

Binoculars Wildlife worksheet Wildlife observation instructions Checklists (optional) Field guides (optional)



3. Explain to the students that each group will take a different measurement, and share their data with each other back in the classroom.

NOTE: If time allows, you can have the groups do more than one of the measurements.

4. Review the sampling instructions for each particular measurement. Have the students fill out the site observations section of the student worksheets before beginning their measurements.

5. Have the students record their results onto the student worksheets. You can choose to have one record keeper per group, or have each student record all the information. You may also want to suggest to your students that they take turns conducting the measurements throughout the process.

ACTIVITY EXTENSIONS:

• Compare your results to results from other streams or other locations on the same stream.

• Sample the same station on multiple dates (fall vs. spring) to compare results.

• Use the activity What's in the Water? to collect chemical measurements in the stream. Have students hypothesize about the links between the riparian zone and the water chemistry. See discussion questions below.

• Map the stream segment – e.g., width of riparian zone, areas with canopy cover, different types of plants.

• Compare (graphically) different reaches of a river – e.g. a more developed area versus a pristine one, such as a city site versus the headwaters.

• Discuss big rivers versus small streams - how do the functions of riparian zones differ?

Further Discussion:

1. How do the greenline measurements help us understand how well a stream bank resists erosion?

Fast moving water can cause banks to erode. The greenline is a measure of how well the plants along the water's edge will help the banks resist erosion (bank stability). We determine stability by calculating the proportion of different vegetation types. These are sedges and rushes, shrubs and trees, grasses, forbs (a non-woody plant that is not a grass) and bare ground. Each vegetation type has a different ability to stabilize the banks due primarily to the depth and density of the roots, and whether they are annuals (die back after one year) or perennials (live through the winter). For example, sedges are perennials that have deep, thick root masses that cling to and stabilize soils, while many annuals have shallow or sparse roots that do not contribute to bank stability.

In the field, look at the vegetation in areas where serious erosion has occurred compared to areas with stable banks. Think about how different land uses affect vegetation and therefore bank stability.

2. How does the canopy cover affect the physical properties of the stream itself?

Canopy cover provides shade and is important in keeping water temperatures low in small "headwater" streams. Many fish and other aquatic organisms are sensitive to high temperatures, and may disappear from streams that have lost their shade.

The canopy of a stream also represents the leaves and debris that may fall directly into the stream. This external input of material is an important source of food and shelter for the fish and other organisms living in these small streams. The relative importance of canopy cover (both for shade and for input of organic material) decreases as a river gets

3. How do humans affect the health of the riparian zone?

The riparian zone is a very small area compared to the entire land area of a watershed, and humans can have a serious impact on this important ecosystem through different types of activities.

• Clearing: Riparian areas are often cleared for agriculture, logging, or housing and other development. This can lead to destabilized banks, heavy erosion and loss of stream and riparian functions.

• Introduced species: Many riparian areas are affected throughout the world by introduced species, which take over the riparian area and radically change the habitat. Species such as russian olive, tamarisk, and purple loosestrife may form "monocultures," replacing native plants and resulting in a serious loss of plant and animal diversity and a loss in other riparian functions such as storing and filtering wastes.

• Grazing: While grazing by cattle and other livestock has been shown to be compatible with healthy riparian areas, the type of grazing is extremely important. Most riparian areas can handle short term, "intensive" grazing, with sufficient recovery time. Continuous grazing in a riparian area can limit the plants' abilities to recover and may ultimately lead to loss of vegetation or a change in species.

• Recreation: Recreationalists flock to riparian areas, but may "love them to death." Trampling, multiple trails, wood removal for campfires, and littering all can impair riparian areas.

4. Why would a stream area be a good habitat for wildlife?

The diversity of plant species (from small annuals to dense sedges to tall trees) provides food and shelter for a wide diversity of animals, ranging from insects to birds to mammals. One function of riparian areas that people sometimes neglect is their use as a "corridor" for wildlife, providing connecting routes across otherwise dry or uninhabitable landscapes. These corridors are important not only for migration, but also for connecting different breeding populations.

Greenline Worksheet

Name:	
Date: _	

Gro	up #:	
Site	ID:	

SITE OBSERVATIONS:

Type of waterbody (e.g., stream, lake, wetland):_____

Water appearance (e.g., rapid moving or slow, still water):_____

What type of land uses are in the immediate area?_____

What type of land uses are in the surrounding area?_____

	Vegetation Categories				
		Rooted ints	Shallow Rooted Plants		
	Sedges & Rushes	Shrubs & Trees	Grasses	Forbs	Bare Ground
Row 1: Record each observation as a slash mark in the appropriate box.					
Row 2: Total number of observations for each category.					
Row 3: Total number of observations for the entire greenline (sum of all observations in Row 2).					
Row 4: Proportion of each category (divide row 2 values by total in row 3).					
Row 5: Multiply each value in row 4 by the factor in each category. Record in row 6	X 10	X 8	X 6	X 3	X 1
Row 6: Score for each category.					

Total Score (add up all scores in Row 6):_____

- Site Scores:
- 7 10 healthy banks
- 4-7 semi healthy banks
- 0-4 unhealthy banks

The higher the score, the more the stream banks will resist erosion.

Worksheet

Canopy Cover

Name:	
Date:	

Group #: _	
Site ID:	

SITE OBSERVATIONS:

Type of waterbody (e.g., stream, lake, wetland):
Water appearance (e.g., clear, brown, foamy, milky):
What type of land uses are in the immediate area?
What type of land uses are in the surrounding area?

	"Miss" (Open sky)	"Hit" (Vegetation)
At each step along the water's edge, record with a slash whether you see a "miss" (open sky) or a "hit" (vegetation) in your ocular tube.		
Total # of slash marks for each category.		
Total number of observations.		
% Canopy cover: Divide total "hits" by total observations and multiply by 100.		

The more "covered" area available, the more shading the stream receives. This keeps the water cool, provides food for aquatic organisms, and woody debris that falls into the stream provides fish habitat.

Ground Cover

4-7 semi healthy banks 0-4 unhealthy banks

	Transects Perpendicular to the Stream (20 paces per transect)					
	1	2	3	4	5	Category Total (percent of total)
Live vegetation						=
Litter						=
Rocks						=
Bare ground						=

Wildlife

Name:	
Date:	

Group #: ______ Site ID: ______

SITE OBSERVATIONS:

Type of waterbody (e.g., stream, lake, wetland):_____

Water appearance (e.g., clear, brown, foamy, milky):_____

What type of land uses are in the immediate area?

What type of land uses are in the surrounding area?_____

COMMON NAME	How is it using the riparian area?	TYPE OF OBSERVATION
e.g., American Dipper	e.g., food, shelter, traveling through	e.g., V, O

Types of Observations

- V = saw animal
- T = track
- S = scat
- B = bones
- O = other sign such as burrow, teeth marks, nests

76


USU Water Quality Extension - Stream Side Science

Time – 30 minutes
Persons – 2
Materials –
Flagging
• Tape measure
 Greenline worksheet
• Plant guide (optional)

1. Measure a 100 foot stretch along the stream. Place a flag near the water at the beginning and end points.

2. Standing at the first flag, note the vegetation type that is closest to the water and record it in row 1 of the worksheet.

3. Take one pace toward the other flag and stop. A pace is a normal stride you would take while walking. Look toward the water and record the vegetation type closest to the water by placing a slash mark in the appropriate box. See the Teacher Resource page for definitons of vegetation types.



4. Repeat these steps until you reach the other flag.

5. Tally the number of slash marks in each box and record this for each category in row 2.

6. Add up all the observations and record that total in row 3.

7. For each vegetation category, divide the number in row 2 by the number in row 3, and record in row 4. This will give the proportion of the greenline that is made up of that vegetation category.

8. For each vegetation category, multiply the number in row 4 by the factor in row 5 and record in row 6. This will give you the "site score" for each vegetation category. Because sedges and rushes have the strongest roots and prevent erosion the best, they receive the highest factor - "10." Bare ground doesn't prevent erosion so it receives the lowest factor – "1."

9. Add the individual site scores in row 6 to get the "total site score" for that stretch of stream.

10. Compare the site score to the "site scores" box on the worksheet to determine the health of the greenline.

Canopy Cover

Гіте – 30 minutes
Persons – 2
Materials –
• Ocular tube
 Measuring tape
 Canopy cover worksheet

1. Measure a 100 foot stretch along the stream. Place a flag near the water at the beginning and end points or use the same measurements set out by the greenline group.

2. Standing at the first flag, point the ocular tube straight into the air (90 degree angle) and look through it with one eye. Your partner who is recording data can tell you how to adjust the tube until it is pointing as straight up as possible.

3. Tell the recorder whether the "X" at the end of the tube points at sky (a "miss") or a part of a tree or bush (a "hit"). Record this in the first row on the canopy cover worksheet.

4. Take one pace toward the other flag and stop. A pace is a normal stride you would take while walking. Again, point the ocular tube straight into the air and record a hit or a miss.

5. Repeat these steps until you reach the other flag.

6. Add the total hits and misses and record in the second row.

7. Add the two scores recorded in row 2. This will tell you the "total number of observations" you took along the transect (the greenline). Record this total in row 3.

8. Divide the number of "hits" in row 2 by the total observations in row 3 and multiply by 100. This will give you the percentage of canopy cover for the transect.

Canopy Cover Transect

Time – 35 minutes Persons - 2

Materials -

• Measuring tape

• Ground cover worksheet

Note: Riparian ground cover transects start at the stream edge and extend 20 paces away from the stream, into the riparian vegetation.

Ground Cover

1. You will collect data along five separate transects in your stream stretch, spaced out at approximately equal distances along your stream reach. If possible, you should run two transects on one side of the stream and three on the other to get a better picture of the total riparian zone. Refer to the figure to the right for help locating these transects.

2. Begin at your first transect. Starting at the stream's edge, take one pace away from the stream. Touch your finger to the ground at the tip of your front foot.

3. Note the ground cover type that your finger touches. The categories are: bare ground, live vegetation, litter (dead vegetation or sticks) or rock. Record the type with a slash in the appropriate box on the ground cover worksheet. Note that each column on the data chart is for a separate transect.

4. Repeat steps 2-3 for 20 paces. Then move on to the second transect. Repeat for all 5 transects.

5. When you've finished with all five transects, add the totals for each row (cover type). This will give you the percentage of each type of ground cover in the riparian zone. To check your math, add your percentages for each ground cover type. They should total 100%.

The percentage of each ground cover type provides a measure of ground cover that can be compared to other sites or used to compare changes over time (between different years or seasons). As a general rule, though, a healthy riparian zone will be covered by a mixture of litter, rock and vegetation. Important exceptions to this are desert streams, which have very sandy banks.

Ground Cover Transect

Wildlife

Time – 35 minutes Persons – 2

Materials –

- Binoculars
- Wildlife worksheet
- Field guides
- Field checklists (optional)
- Magnifying glasses (optional)

Note: Wildlife and birds are shy and may be hard to observe. You may want to send your wildlife observers directly upstream or downstream of your transect so the noise and activity of the vegetation survey won't interfere with their observations. Only do this if it is safe and practical.

1. Slowly walk a 100 foot length along the stream's edge. Look carefully for tracks, scat

or other animal signs. Also watch for birds, mammals, reptiles, or amphibians as you

walk. Record all species that can be identified by sight or sound on the wildlife

worksheet. Also note how the wildlife may have been using the riparian area, and the

type of oberservation.

2. Return to approximately the middle of the length of stream you just walked, and walk

20 paces away from the stream's edge. Stand quietly, watching and listening for two

minutes. Record all species that can be identified.

3. Slowly walk back toward the stream, looking for tracks, scat or other animal signs.

Once you have reached the stream, listen and watch for another two minutes. Repeat this procedure at three points (more if time allows) along the stream.

4. After sampling, spend any extra time investigating the entire site looking for animals or signs of them.

How to make an ocular tube

3. Note the ground cover type that your finger touches. The categories are: bare ground,

live vegetation, litter (dead vegetation or sticks) or rock. Record the type with a slash in the appropriate box on the ground cover worksheet. Note that each column on the data chart is for a separate transect.

4. Repeat steps 2 - 3 for 20 paces. Then move on to the second transect. Repeat for all 5 transects.

5. When you've finished with all five transects, add the totals for each row (cover type). This will give you the percentage of each type of ground cover in the riparian zone. To check your math, add your percentages for each ground cover type. They should total 100%. An ocular tube is used to sample canopy cover. It is a very simple device, but it removes bias from field sampling canopy cover.




Illustration: Holly Broome-Hyer

Materials

- 6 in. of 1 in. wide PVC or other material such as a cardboard tube.
- 2 paper clips
- Duct tape

Directions

1. Cut a 6 in. length of tube.

2. Make four notches every 90 degrees on one end.

3. Straighten paper clips and lay into notches. Bend excess length over the outside of the tube.

4. Duct tape around the ends of the paper clips to hold them in place. Use duct tape over the edges of the viewing end of the tube to make a smooth surface.

Field Checklist for

Mammals

\checkmark	Common Name	Scientific Name
	Masked Shrew	Sorex cinereus
	Dusky Shrew	Sorex obscurus
	Northern Water Shrew	Sorex palustris
	Black Bear	Ursus americanus
	Raccoon	Procyon lotor
	Longtail Weasel	Mustela frenata
	Mink	Mustela vison
	River Otter	Lutra canadensis
	Striped Skunk	Mephitis mephitis
	Coyote	Canis latrans
	Red Fox	Vulpes fulpa
	Mountain Lion	Felis concolor
	Bobcat	Lynx rufus
	Yellowbelly Marmot	Marmota flaveventris
	Northern Pocket Gopher	Thomomys talpoides
	Beaver	Castor canadensis
	Meadow Vole	Microtus pennslvanicus
	Muskrat	Ondantra zibethica
	Porcupine	Erethizon dorsatum
	Mountain Cottontail	Sylvilagus nutalli
	Elk	Cervus elaphus
	Mule Deer	Odocoileus hemionus
	Moose	Alces alces
	Pronghorn	Antilocapra Americana
	Douglas' Squirrel	Tamiasciurus douglasii

82


USU Water Quality Extension - Stream Side Science

Field Checklist for Birds

\checkmark	Riparian Species of Utah * = southern Utah species	
	Ring-necked Pheasant	
	Wild Turkey	
	Great Blue Heron	
	Cooper's Hawk	
	Red-tailed Hawk	
	American Kestrel	
	Yellow-billed Cuckoo	
	Western Screech-owl	
	Great Horned Owl	
	Black-chinned Hummingbird	
	Belted Kingfisher	
	Downy Woodpecker	
	Willow Flycatcher	
	Cassin's Kingbird	
	Eastern Kingbird	
	Bell's Vireo *	
	Plumbeous Vireo	
	Brown Creeper	
	Bewick's Wren	

House Wren
American Dipper
Swainson's Thrush
American Robin
Gray Catbird
European Starling
Cedar Waxwing
Yellow Warbler
Wilson's Warbler
Yellow-breasted Chat
Spotted Towhee
Abert's Towhee *
Fox Sparrow
Song Sparrow
Lincoln's Sparrow
Black-headed Grosbeak
Blue Grosbeak
Lazuli Bunting
Brown-headed Cowbird
Bullock's Oriole
Lesser Goldfinch *

Plant Identification



Vegetation Types

Grasses – These have hollow stems that are jointed and leaves with parallel veins. The leaves come off the stem in opposite directions.

Grasslikes sedges – These resemble grasses but have solid, triangular stems with no joints. The leaves have parallel veins but they come off the stem in three directions. This group also includes rushes which have round, hollow stems with very small or no leaves. **Forbs** – These generally have broad leaves with net-like veins. The stems are solid or spongy and they die back to the ground every year.

Shrubs and trees – These have woody stems that remain alive all year. The leaves tend to have net-like veins. Rarely do shrubs grow taller than 13 feet. Trees are similar to shrubs in that they generally have a single woody stem but grow taller than 13 feet.

This page reprinted with permission from the Utah Stream Team Manual

Notes

USU Water Quality Extension - Stream Side Science

